

Statistical Methods for Data Science

Lessons 26 - Confidence intervals: mean, proportion, linear regression

Salvatore Ruggieri

Department of Computer Science
University of Pisa
salvatore.ruggieri@unipi.it

From point estimate to interval estimate

Estimator and point estimate

A *statistics* is a function of $h(X_1, \dots, X_n)$ of r.v.'s.

An *estimator* of a parameter θ is a statistics $T_n = h(X_1, \dots, X_n)$ intended to provide information about θ .

A *point estimate* t of θ is $t = h(x_1, \dots, x_n)$ over realizations of X_1, \dots, X_n .

- Sometimes, a *range* of plausible values for an unknown parameter is preferred
- Idea: *confidence interval* is an interval for which we can be confident the unknown parameter is in with a specified probability (*confidence level*)

Example

- From the Chebyshev's inequality:

$$P(|Y - \mu| < k\sigma) \geq 1 - \frac{1}{k^2}$$

For $Y = \bar{X}_n$, $k = 2$ and $\sigma = 100$ Km/s:

$$P(|\bar{X}_n - \mu| < 200) \geq 0.75$$

- ▶ i.e., $\bar{X}_n \in (\mu - 200, \mu + 200)$ with probability $\geq 75\%$ *[random variable in a fixed interval]*
 - ▶ or, $\mu \in (\bar{X}_n - 200, \bar{X}_n + 200)$ with probability $\geq 75\%$ *[fixed value in a random interval]*
- $(\bar{X}_n - 200, \bar{X}_n + 200)$ is an interval estimator of the unknown μ
 - ▶ the interval contains μ with probability $\geq 75\%$
- Let $t = 299852.4$ be the point estimate (realization of $T = \bar{X}_n$)
- $\mu \in (t - 200, t + 200) = (299652.4, 300052.4)$ is correct with confidence $\geq 75\%$

Table 17.1. Michelson data on the speed of light.

850	740	900	1070	930	850	950	980	980	880
1000	980	930	650	760	810	1000	1000	960	960
960	940	960	940	880	800	850	880	900	840
830	790	810	880	880	830	800	790	760	800
880	880	880	860	720	720	620	860	970	950
880	910	850	870	840	840	850	840	840	840
890	810	810	820	800	770	760	740	750	760
910	920	890	860	880	720	840	850	850	780
890	840	780	810	760	810	790	810	820	850
870	870	810	740	810	940	950	800	810	870

The better the estimator the better (smaller) the intervals

- Assume $X_i \sim N(\mu, \sigma^2)$. Hence, $\bar{X}_n \sim N(\mu, \sigma^2/n)$ and:

$$Z_n = \sqrt{n} \frac{\bar{X}_n - \mu}{\sigma} \sim N(0, 1)$$

- $P(|Z_n| \leq 1.15) = P(-1.15 \leq Z_n \leq 1.15) = \Phi(1.15) - \Phi(-1.15) = 0.75$
 - $-1.15 = q_{0.125}$ and $1.15 = q_{0.875}$ are called *the critical values* for achieving 75% probability
- Going back to \bar{X}_n :

$$P(-1.15 \leq \sqrt{n} \frac{\bar{X}_n - \mu}{\sigma} \leq 1.15) = P(\bar{X}_n - 1.15 \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{X}_n + 1.15 \frac{\sigma}{\sqrt{n}}) = 0.75$$

- $\mu \in (t - 1.15 \frac{200}{\sqrt{100}}, t + 1.15 \frac{200}{\sqrt{100}}) = \underline{(t - 23, t + 23)}$ is correct with confidence = 75%

Confidence intervals

CONFIDENCE INTERVALS. Suppose a dataset x_1, \dots, x_n is given, modeled as realization of random variables X_1, \dots, X_n . Let θ be the parameter of interest, and γ a number between 0 and 1. If there exist sample statistics $L_n = g(X_1, \dots, X_n)$ and $U_n = h(X_1, \dots, X_n)$ such that

$$P(L_n < \theta < U_n) = \gamma$$

for every value of θ , then

$$(l_n, u_n),$$

where $l_n = g(x_1, \dots, x_n)$ and $u_n = h(x_1, \dots, x_n)$, is called a $100\gamma\%$ confidence interval for θ . The number γ is called the *confidence level*.

- Sometimes, only have $P(L_n < \theta < U_n) \geq \gamma$ [*conservative* $100\gamma\%$ confidence interval]
 - ▶ E.g., the interval found using Chebyshev's inequality
- There is no way of knowing if $l_n < \theta < u_n$ (interval is correct)
- We only know that we have probability γ of covering θ
- Notation: $\gamma = 1 - \alpha$ where α is called the *significance level*
 - ▶ E.g., 95% confidence level equivalent to 0.05 significance level

Seeing theory simulation

Confidence interval for the mean

- Let X_1, \dots, X_n be a random sample and $\mu = E[X_i]$ to be estimated
- Problem: confidence intervals for μ ?
 - ▶ Normal data
 - with known variance
 - with unknown variance
 - ▶ General data (with unknown variance)
 - large sample, i.e., large n
 - bootstrap (next lesson)

Critical values

Critical value

The (right) *critical value* z_p of $Z \sim N(0, 1)$ is the number with right tail probability p :

$$P(Z \geq z_p) = p$$


- Alternatively, $p = 1 - \Phi(z_p) = 1 - P(Z \leq z_p)$.
 - ▶ This is why Table B.1 of the textbook is given for $1 - \Phi()$
- Alternatively, $\Phi(z_p) = 1 - p$, i.e., z_p is the $(1 - p)$ th quantile
- By $P(Z \geq z_p) = P(Z \leq -z_p) = p$, we have:

$$P(Z \geq -z_p) = 1 - P(Z \leq -z_p) = 1 - p$$

and then:

$$z_{1-p} = -z_p$$

- ▶ E.g., $z_{0.975} = -z_{0.025} = -1.96$ and $z_{0.025} = -z_{0.975} = 1.96$



CI for the mean: normal data with known variance

- Dataset x_1, \dots, x_n realization of random sample $X_1, \dots, X_n \sim N(\mu, \sigma^2)$
- Estimator $\bar{X}_n \sim N(\mu, \sigma^2/n)$ and the scaled mean:

$$Z = \sqrt{n} \frac{\bar{X}_n - \mu}{\sigma} \sim N(0, 1) \quad (1)$$

- Confidence interval for Z :

$$P(c_l \leq Z \leq c_u) = \gamma \quad \text{or} \quad P(Z \leq c_l) + P(Z \geq c_u) = \alpha = 1 - \gamma$$

- Symmetric split:

$$P(Z \leq c_l) = P(Z \geq c_u) = \alpha/2$$

Hence $c_u = -c_l = z_{\alpha/2}$, and by (1):

$$P\left(\bar{X}_n - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \bar{X}_n + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}\right) = 1 - \alpha = \gamma$$

$(\bar{x}_n - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}, \bar{x}_n + z_{\alpha/2} \frac{\sigma}{\sqrt{n}})$ is a $100\gamma\%$ or $100(1 - \alpha)\%$ confidence interval for μ

One-sided confidence intervals

- One-sided confidence intervals (*greater-than*):

$$P(L_n < \theta) = \gamma$$

Then (l_n, ∞) is a $100\gamma\%$ or $100(1 - \alpha)\%$ one-sided confidence interval

- l_n is called the *lower confidence bound*
- Normal data with known variance:

$$P(\bar{X}_n - z_\alpha \frac{\sigma}{\sqrt{n}} \leq \mu) = 1 - \alpha = \gamma$$

$(\bar{x}_n - z_\alpha \frac{\sigma}{\sqrt{n}}, \infty)$ is a $100\gamma\%$ or $100(1 - \alpha)\%$ one-sided confidence interval for μ

See R script

CI for the mean: normal data with unknown variance

- Use the unbiased estimator of σ^2 and its estimate

$$S_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2 \qquad s_n^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}_n)^2$$

- ▶ and then S_n^2/n is an unbiased estimator of $\text{Var}(\bar{X}_n) = \sigma^2/n$
- The following transformation is called *studentized mean*: $T = \sqrt{n} \frac{\bar{X}_n - \mu}{S_n} \sim t(n-1)$

DEFINITION. A continuous random variable has a *t-distribution with parameter* m , where $m \geq 1$ is an integer, if its probability density is given by

$$f(x) = k_m \left(1 + \frac{x^2}{m}\right)^{-\frac{m+1}{2}} \quad \text{for } -\infty < x < \infty,$$

where $k_m = \Gamma(\frac{m+1}{2}) / (\Gamma(\frac{m}{2}) \sqrt{m\pi})$. This distribution is denoted by $t(m)$ and is referred to as the *t-distribution with m degrees of freedom*.

- Student t-distribution $X \sim t(m)$:
 - ▶ for $m = 1$, it is the Cauchy distribution
 - ▶ $E[X] = 0$ for $m \geq 2$, and $\text{Var}(X) = m/(m-2)$ for $m \geq 3$
 - ▶ For $m \rightarrow \infty$, $X \rightarrow N(0, 1)$

Some history on its discovery

See R script

CI for the mean: normal data with unknown variance

- Dataset x_1, \dots, x_n realization of random sample $X_1, \dots, X_n \sim N(\mu, \sigma^2)$

Critical value

The (right) *critical value* $t_{m,p}$ of $T \sim t(m)$ is the number with right tail probability p :

$$P(T \geq t_{m,p}) = p$$

- Same properties of z_p
- From the studentized mean:

$$T = \sqrt{n} \frac{\bar{X}_n - \mu}{S_n} \sim t(n-1)$$

to confidence interval:

$$P\left(\bar{X}_n - t_{n-1,\alpha/2} \frac{S_n}{\sqrt{n}} \leq \mu \leq \bar{X}_n + t_{n-1,\alpha/2} \frac{S_n}{\sqrt{n}}\right) = 1 - \alpha = \gamma$$

$\left(\bar{X}_n - t_{n-1,\alpha/2} \frac{S_n}{\sqrt{n}}, \bar{X}_n + t_{n-1,\alpha/2} \frac{S_n}{\sqrt{n}}\right)$ is a $100\gamma\%$ or $100(1 - \alpha)\%$ confidence interval for μ

See R script

CI for the mean: general data with unknown variance

- Dataset x_1, \dots, x_n realization of random sample $X_1, \dots, X_n \sim F$
- A variant of CLT states that for $n \rightarrow \infty$

$$T = \sqrt{n} \frac{\bar{X}_n - \mu}{S_n} \rightarrow N(0, 1)$$

- For large n , we make the approximation:

[how large should n be?]

$$T = \sqrt{n} \frac{\bar{X}_n - \mu}{S_n} \approx N(0, 1)$$

and then

$$P\left(\bar{X}_n - z_{\alpha/2} \frac{S_n}{\sqrt{n}} \leq \mu \leq \bar{X}_n + z_{\alpha/2} \frac{S_n}{\sqrt{n}}\right) \approx 1 - \alpha = \gamma$$

$(\bar{X}_n - z_{\alpha/2} \frac{S_n}{\sqrt{n}}, \bar{X}_n + z_{\alpha/2} \frac{S_n}{\sqrt{n}})$ is a $100\gamma\%$ or $100(1 - \alpha)\%$ confidence interval for μ

See R script

General form of Wald confidence intervals

$$\theta \in \hat{\theta} \pm z_{\alpha/2} \text{se}(\hat{\theta}) \quad \text{or} \quad \theta \in \hat{\theta} \pm t_{\alpha/2} \text{se}(\hat{\theta})$$

- They originate from the **Wald test statistics**:

$$T = \frac{\hat{\theta} - \theta}{\sqrt{\text{Var}(\hat{\theta})}} = \frac{\hat{\theta} - \theta}{\text{se}(\hat{\theta})}$$

- Importance of standard error $\text{se}(\hat{\theta})$ of estimators!
- Limitation: asymptotic, symmetric intervals

CI for proportions (e.g., misclassification error)

- Dataset x_1, \dots, x_n realization of random sample $X_1, \dots, X_n \sim \text{Ber}(p)$
 - ▶ $x_i = \mathbb{1}_{y_{\theta}^+(w_i) \neq c_i}$ is 1 for incorrect classification, 0 for correct classification [over a test set]
 - ▶ p is the (unknown) misclassification error of classifier $y_{\theta}^+(\cdot)$
- $B = \sum_{i=1}^n X_i \sim \text{Bin}(n, p)$ and $b = \sum_{i=1}^n x_i$ (number of observed successes)
 - ▶ For small n , we can build exact bounds (p_L, p_U) such that: [Exact interval]

$$l_B = \min_{\theta} \left\{ \sum_{x=B}^n \binom{n}{x} \theta^x (1-\theta)^{n-x} \geq \alpha/2 \right\} \quad u_B = \max_{\theta} \left\{ \sum_{x=0}^B \binom{n}{x} \theta^x (1-\theta)^{n-x} \geq \alpha/2 \right\}$$

- l_B is the smallest θ for which $P(X \geq b) \geq \alpha/2$ for $X \sim \text{Bin}(n, \theta)$
- u_B is the greatest θ for which $P(X \leq b) \geq \alpha/2$ for $X \sim \text{Bin}(n, \theta)$

$$P(l_B \leq p \leq u_B) = 1 - \alpha$$

and then (l_b, u_b) is a $100\gamma\%$ or $100(1 - \alpha)\%$ confidence interval for p

- ▶ Closed form given using quantiles of the F-distribution

CI for proportions (e.g., misclassification error)

- Dataset x_1, \dots, x_n realization of random sample $X_1, \dots, X_n \sim \text{Ber}(p)$
 - ▶ $x_i = \mathbb{1}_{y_{\theta}^+(w_i) \neq c_i}$ is 1 for incorrect classification, 0 for correct classification *[over a test set]*
 - ▶ p is the (unknown) misclassification error of classifier $y_{\theta}^+(\cdot)$
- $B = \sum_{i=1}^n X_i \sim \text{Bin}(n, p)$
 - ▶ For large n , $\text{Bin}(n, p) \approx N(np, np(1-p))$ for $0 \ll p \ll 1$ **[De Moivre–Laplace]**
 - $\text{se}(B) = \sqrt{n\bar{X}_n(1-\bar{X}_n)}$ because $B \sim \text{Bin}(n, p)$
 - We make the approximation: $T = (B - np)/\text{se}(B) \approx N(0, 1)$ and then
$$P\left(\bar{X}_n - z_{\alpha/2} \sqrt{\frac{\bar{X}_n(1-\bar{X}_n)}{n}} \leq p \leq \bar{X}_n + z_{\alpha/2} \sqrt{\frac{\bar{X}_n(1-\bar{X}_n)}{n}}\right) \approx 1 - \alpha = \gamma$$

$(\bar{x}_n - z_{\alpha/2} \sqrt{\frac{\bar{x}_n(1-\bar{x}_n)}{n}}, \bar{x}_n + z_{\alpha/2} \sqrt{\frac{\bar{x}_n(1-\bar{x}_n)}{n}})$ is a $100\gamma\%$ or $100(1-\alpha)\%$ confidence interval for p
 - This is a Wald confidence interval!
- ▶ Drawbacks: symmetric, large sample, skewness, etc. **[Wilson score interval]**

See R script

Determining the sample size

- For a fixed α , the narrower the CI the better (smaller variability)
- Sometimes, we start with an accuracy requirement (maximal width w of the interval):
 - ▶ find a $100(1 - \alpha)\%$ CI (l_n, u_n) such that $u_n - l_n \leq w$
- How to set n to satisfy the w bound?
- Case: normal data with known variance σ^2
 - ▶ CI is $(\bar{X}_n - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}, \bar{X}_n + z_{\alpha/2} \frac{\sigma}{\sqrt{n}})$
 - ▶ Bound on the CI is:

$$2z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq w$$

leading to:

$$n \geq \left(2z_{\alpha/2} \frac{\sigma}{w}\right)^2$$

- Case σ^2 unknown: use estimate $s_n^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}_n)^2$

Confidence intervals for simple linear regression coefficients

- Simple linear regression: $Y_i = \alpha + \beta x_i + U_i$ with $\underline{U_i \sim \mathcal{N}(0, \sigma^2)}$ and $i = 1, \dots, n$
- We have $\hat{\beta} \sim \mathcal{N}(\beta, \text{Var}(\hat{\beta}))$ where $\text{Var}(\hat{\beta}) = \sigma^2/SXX$ is unknown *[see Lesson 20]*
- The Wald statistics is $t(n-2)$ -distributed: *[proof omitted]*

$$\frac{\hat{\beta} - \beta}{\sqrt{\text{Var}(\hat{\beta})}} \sim t(n-2)$$

- For $\gamma = 0.95$:

$$P(-t_{n-2,0.025} \leq \frac{\hat{\beta} - \beta}{\sqrt{\text{Var}(\hat{\beta})}} \leq t_{n-2,0.025}) = 0.95$$

and then a 95% confidence interval is: $\hat{\beta} \pm t_{n-2,0.025} \text{se}(\hat{\beta})$ where $\text{se}(\hat{\beta}) = \hat{\sigma}/\sqrt{SXX}$

- Similarly, we get for α , $\hat{\alpha} \pm t_{n-2,0.025} \text{se}(\hat{\alpha})$

See R script

Confidence intervals of fitted values

- Simple linear regression: $Y_i = \alpha + \beta x_i + U_i$ with $\underline{U_i \sim \mathcal{N}(0, \sigma^2)}$ and $i = 1, \dots, n$
- For the fitted values $\hat{y} = \hat{\alpha} + \hat{\beta}x_0$ at x_0 , a 95% confidence interval is:

$$\hat{y} \pm t_{n-2,0.025} \text{se}(\hat{y})$$

$$\text{where } \text{se}(\hat{y}) = \hat{\sigma} \sqrt{\left(\frac{1}{n} + \frac{(\bar{x}_n - x_0)^2}{SXX}\right)}$$

[see Lesson 21]

- This interval concerns *the expectation of fitted values* at x_0 .
 - ▶ E.g., the mean of predicted values at x_0 is in $[\hat{y} + t_{n-2,0.025} \text{se}(\hat{y}), \hat{y} - t_{n-2,0.025} \text{se}(\hat{y})]$

See R script

Prediction intervals of fitted values

- Simple linear regression: $Y_i = \alpha + \beta x_i + U_i$ with $\underline{U_i \sim \mathcal{N}(0, \sigma^2)}$ and $i = 1, \dots, n$
- For a given *single prediction*, we must also account for the error term U in:

$$\hat{V} = \hat{\alpha} + \hat{\beta}x_0 + U$$

- Assuming $U \sim \mathcal{N}(0, \sigma^2)$, we have $\text{Var}(\hat{V}) = \sigma^2(1 + \frac{1}{n} + \frac{(\bar{x}_n - x_0)^2}{SXX})$
- A 95% confidence interval is:

$$\hat{y} \pm t_{n-2, 0.025} \text{se}(\hat{v})$$

$$\text{where } \text{se}(\hat{v}) = \hat{\sigma} \sqrt{1 + \frac{1}{n} + \frac{(\bar{x}_n - x_0)^2}{SXX}}$$

- A predicted value at x_0 is in $[\hat{y} - t_{n-2, 0.025} \text{se}(\hat{v}) \text{ and } \hat{y} + t_{n-2, 0.025} \text{se}(\hat{v})]$

See R script

Optional reference

- On confidence intervals and statistical tests (with R code)

 Myles Hollander, Douglas A. Wolfe, and Eric Chicken (2014)
Nonparametric Statistical Methods.
3rd edition, *John Wiley & Sons, Inc.*